

THE PION FORM FACTOR: SUDAKOV SUPPRESSIONS AND INTRINSIC TRANSVERSE MOMENTUM

R. Jakob ^{1 2}

Fachbereich Physik, Universität Wuppertal, Gaußstrasse 20,
Postfach 10 01 27, D-5600 Wuppertal 1, Germany

and

P. Kroll ³

Theory Division, CERN, CH-1211 Geneva 23, Switzerland

Abstract

It is demonstrated that any attempt to calculate the perturbative QCD contribution to the pion form factor requires the inclusion of intrinsic transverse momentum besides Sudakov form factors. For momentum transfers of the order of a few GeV the intrinsic transverse momentum leads to a substantial suppression of the perturbative QCD contribution.

¹Supported by the Deutsche Forschungsgemeinschaft

²E-mail: rainer@wpts0.physik.uni-wuppertal.de

³On leave of absence from Fachbereich Physik, Universität Wuppertal, Germany

There is general agreement that perturbative QCD in the framework of the hard-scattering picture [1] is the correct description of exclusive reactions at asymptotically large momentum transfer. However, the applicability of this approach at experimentally accessible momentum transfers, typically a few GeV, was questioned [2, 3]. It was asserted that in the few GeV region the hard-scattering picture accumulates strong contributions from the soft end-point regions, rendering the perturbation calculation inconsistent. Recently, this statement was disproved by Sterman and collaborators [4, 5, 6]. The essential point of their work is that the customarily neglected transverse momenta of the quarks as well as the Sudakov corrections are taken into account. Because of that the perturbative QCD contribution becomes self-consistent, even for momentum transfers as low as a few GeV, and seem to dominate form factors and cross-sections. For the electromagnetic form factor of the pion, the simplest exclusive quantity, it has been shown [5] that perturbative QCD can readily be applied for momentum transfers larger than $20\Lambda_{QCD}$. It should be noted that the analyses of [4, 5, 6] involve no phenomenological parameter.

However, it seems to us that an important element is missed in the analyses of [5, 6], namely the intrinsic transverse momentum dependence of the hadronic wave function. As we are going to demonstrate for the case of the pion form factor, the inclusion of that $k_\perp$ -dependence substantially suppresses the contribution from perturbative QCD. Although self-consistently calculated, this contribution then seems to be too small with respect to the data. We therefore suspect that there are sizable soft contributions to the pion form factor in the few GeV region, a conjecture already expressed in [2, 3].

In order to make our point of view clear it is useful to repeat the essential steps in the derivation of the hard-scattering formula for the pion electromagnetic form factor [1]. The starting point is the Drell-Yan formula [7] in which the pion form factor is expressed as an overlap of the initial- and final-state light-cone wave functions

$$F_\pi(Q^2) = \int \frac{dx_1 d^2k_\perp}{16\pi^3} \Psi^*(x_1, \vec{k}_\perp + x_2 \vec{q}) \Psi(x_1, \vec{k}_\perp), \quad (1)$$

where $Q^2 = \vec{q}^2$ is the momentum transfer, x_1 is the usual longitudinal momentum fraction carried by the quark and $\vec{k}_\perp$ its transverse momentum with respect to its parent hadron. The momentum of the antiquark is characterized by $x_2 = 1 - x_1$ and $-\vec{k}_\perp$; Ψ is the wave function of the valence Fock state ⁴. It satisfies the normalization condition

$$\int \frac{dx_1 d^2k_\perp}{16\pi^3} |\Psi(x_1, \vec{k}_\perp)|^2 = P_{q\bar{q}} \leq 1. \quad (2)$$

Contributions from higher Fock states are neglected in (1) since, at large momentum transfer, they are suppressed by inverse powers of Q^2 . As the inspection of (1) reveals, the large Q behaviour of the pion form factor is controlled by the tail of the wave function at large $\vec{k}_\perp$. The crucial point is that the tail of the wave function can be calculated, within perturbative QCD, from the soft part of the wave function by means of a homogeneous Bethe-Salpeter equation. Absorbing the perturbative kernel of the Bethe-Salpeter equation in a hard-scattering amplitude T_H , expression (1) can be converted into

$$F_\pi(Q^2) = \int \frac{dx_1 d^2k_\perp}{16\pi^3} \int \frac{dy_1 d^2l_\perp}{16\pi^3} \Psi_0^*(y_1, \vec{l}_\perp) T_H(x_1, y_1, Q, \vec{k}_\perp, \vec{l}_\perp) \Psi_0(x_1, \vec{k}_\perp). \quad (3)$$

Now Ψ_0 represents the soft part of the pion wave function, i.e. the full wave function with the perturbative tail removed from it. The perturbative tail of Ψ behaves as $1/k_\perp^2$ for large $k_\perp$

⁴Strictly speaking Ψ is the $S = 0, L_z = 0$ component of the valence Fock state wave function.

whereas Ψ_0 vanishes as $\sim 1/k_\perp^4$ or faster. To lowest order in perturbation theory, the hard-scattering amplitude is to be calculated from the one-gluon-exchange diagrams. Keeping the transverse momenta of the quarks, $\vec{k}_\perp$ and $\vec{l}_\perp$, only in the gluon propagators, where they matter most ⁵, the hard-scattering amplitude reads

$$T_H(x_1, y_1, Q, \vec{k}_\perp, \vec{l}_\perp) = \frac{16\pi \alpha_s(\mu) C_F}{x_1 y_1 Q^2 + (\vec{k}_\perp + \vec{l}_\perp)^2}, \quad (4)$$

where we have made use of the symmetry of the pion wave function under the replacement $x_1 \leftrightarrow x_2$ (C invariance). In eq. (4) α_s is the usual strong coupling constant to be evaluated at a renormalization scale μ and $C_F (= 4/3)$ is the colour factor.

At large Q one may neglect the $k_\perp$ - and $l_\perp$ -dependence in the gluon propagator as well; T_H can then be pulled out of the transverse momentum integrals, and these integrations apply only to the wave functions. Defining distribution amplitudes (DA) by

$$\frac{f_\pi}{2\sqrt{6}} \phi(x_1, \mu) = \int \frac{d^2 k_\perp}{16\pi^3} \Psi_0(x_1, \vec{k}_\perp), \quad (5)$$

one arrives at the celebrated hard-scattering formula for the pion form factor

$$F_\pi^{HSP}(Q^2) = \frac{f_\pi^2}{24} \int dx_1 dy_1 \phi^*(y_1, \mu) T_H(x_1, y_1, Q, \mu) \phi(x_1, \mu), \quad (6)$$

which is valid for $Q \rightarrow \infty$.

The DA is defined such that

$$\int_0^1 dx_1 \phi(x_1, \mu) = 1. \quad (7)$$

An immediate consequence of the definitions (5) and (7) is that the constraint from the $\pi \rightarrow \mu\nu$ decay [8] is automatically satisfied:

$$\frac{f_\pi}{2\sqrt{6}} = \int \frac{dx_1 d^2 k_\perp}{16\pi^3} \Psi_0(x_1, \vec{k}_\perp), \quad (8)$$

where $f_\pi (= 133 \text{ MeV})$ is the usual π decay constant. The integral in (5) has to be cut off at a scale of order Q , which leads to a very mild dependence of the DA on the renormalization scale (QCD evolution). For most applications of perturbative QCD to exclusive reactions, the μ -dependence of the DAs can be ignored since one is only interested in a rather limited region of Q .

The neglect of the transverse momentum dependence in (4) is a bad approximation in the end-point regions where one of the momentum fractions, x_i or y_i , $i = 1, 2$, tends to zero. This approximation is after all responsible for the theoretical inconsistencies mentioned in the introduction. Li and Sterman [5] have suggested to retain the transverse momentum dependence of T_H . Moreover, Sudakov corrections, suppressing the contributions from the dangerous soft regions, ought to be taken into account. In order to include these corrections it is advantageous to reexpress eq. (3) in terms of the Fourier transform variable $\vec{b}$ in the transverse configuration space ⁶

$$F_\pi^{pert}(Q^2) = \int_0^1 \frac{dx_1 dy_1}{(4\pi)^2} \int_{-\infty}^{\infty} d^2 b \hat{\Psi}_0^*(y_1, \vec{b}, w) \hat{T}_H(x_1, y_1, Q, b, t) \hat{\Psi}_0(x_1, -\vec{b}, w) \times \exp[-S(x_1, y_1, Q, b, t)] \quad (9)$$

⁵The effect of the $k_\perp$ -dependence in the fermion propagators has been investigated by Li [6]; it leads to a reduction of the results for F_π by about 10% for $Q > 3 \text{ GeV}$.

⁶Since T_H depends only on $\vec{k}_\perp + \vec{l}_\perp$ there is merely one independent variable in the transverse configuration space.

where the Fourier transform of a function $f = f(\vec{k}_\perp)$ is defined by

$$\hat{f}(\vec{b}) = \frac{1}{(2\pi)^2} \int d^2 k_\perp e^{-i\vec{b} \cdot \vec{k}_\perp} f(\vec{k}_\perp). \quad (10)$$

The Fourier-transformed hard-scattering amplitude reads

$$\hat{T}_H(x_1, y_1, Q, b, t) = 8 \alpha_s(t) C_F K_0(\sqrt{x_1 y_1} Q b), \quad (11)$$

where K_0 is the modified Bessel function of order zero and t is the largest mass scale appearing in $\hat{T}_H$:

$$t = \text{Max}(\sqrt{x_1 y_1} Q, w = 1/b). \quad (12)$$

The factor $\exp[-S]$ in (9), termed the Sudakov factor, comprises the radiative corrections. The function S is given by [4]

$$S(x_1, y_1, Q, b, t) = \sum_{i=1}^2 [s(x_i, Q, b) + s(y_i, Q, b)] - \frac{2}{\beta_1} \ln \frac{\ln(t/\Lambda_{QCD})}{\ln(1/b\Lambda_{QCD})}. \quad (13)$$

The last term in (13) arises from the application of the renormalization group. The lengthy expression for the Sudakov exponent s , which includes all leading and next-to-leading logarithms, is given explicitly in [5]. The most important term in it is the double logarithm

$$\frac{2}{3\beta_1} \ln \frac{\xi Q}{\sqrt{2}\Lambda_{QCD}} \ln \frac{\ln(\xi Q/\sqrt{2}\Lambda_{QCD})}{\ln(1/b\Lambda_{QCD})}, \quad (14)$$

where ξ is one of the fractions, x_i or y_i , and $\beta_1 = (33 - 2n_f)/12$. For small b , i.e. at small transverse separation of quark and antiquark, there is no suppression from the Sudakov factor⁷. As b increases the Sudakov factor decreases, reaching zero at $b = 1/\Lambda_{QCD}$. For b larger than $1/\Lambda_{QCD}$ the Sudakov factor is set to zero. Owing to this cut-off the singularity of α_s is avoided without introducing a phenomenological parameter (e.g. a gluon mass). For $Q \rightarrow \infty$ the Sudakov factor dumps any contribution except those from configurations with small quark-antiquark separation. In other words, the hard-scattering contributions dominate the pion form factor asymptotically.

Li and Sterman have explored the improved hard-scattering formula (9) on the basis of customary wave functions, neglecting the QCD evolution and the intrinsic transverse momentum dependence of the wave functions. Their numerical studies have revealed that the modified perturbative approach is self-consistent for $Q > 20\Lambda_{QCD}$ in the sense that less than, say, 50% of the result is generated by soft gluon exchange ($\alpha_s > 0.7$). In the few GeV region the values for F_π as obtained by Li and Sterman are somewhat smaller than those provided by the hard-scattering formula (6) and are, perhaps, smaller than the experimental values [9]. The data may however suffer from large systematic errors [10]. This prevents us from giving a definite conclusion about the agreement or disagreement between data and theory.

The approach proposed by Sterman and collaborators [4, 5, 6] certainly constitutes an enormous progress in our understanding of exclusive reactions at large momentum transfer. We believe, however, that in any practical application of that approach one has to allow for an intrinsic transverse momentum dependence of the hadronic wave function, although, admittedly, this requires a new phenomenological element in the calculation. In order to demonstrate the

⁷Actually the Sudakov factor includes a small enhancement in the low- b region. Li and Sterman [5] neglect this enhancement and set any factor $\exp[-s(\xi, b, Q)]$ to unity in this region and whenever $\xi < \sqrt{2}/bQ$. We apply this recipe too.

importance of the intrinsic transverse momentum for the case of the pion form factor, we need realistic pion wave functions.

The pion wave function: In accordance with (5), (7) and (8) we write the soft part of the valence quark Fock state wave function as

$$\Psi_0(x_1, \vec{k}_\perp) = \frac{f_\pi}{2\sqrt{6}} \phi(x_1) \Sigma(x_1, \vec{k}_\perp), \quad (15)$$

the function Σ being normalized in such a way that

$$\int \frac{d^2 k_\perp}{16\pi^3} \Sigma(x_1, \vec{k}_\perp) = 1. \quad (16)$$

The probability of the valence quark Fock state is given by (2)⁸. A further constraint of the wave function comes from the charge radius of the pion. The (transverse) radius of the valence Fock state should be smaller than (or at best equal to) the full radius. Or, in terms of transverse momentum, we have a lower limit on the mean square transverse momentum that is defined by

$$\langle k_\perp^2 \rangle = \frac{f_\pi^2}{24 P_{q\bar{q}}} \int \frac{dx_1 d^2 k_\perp}{16\pi^3} k_\perp^2 |\phi(x_1)|^2 |\Sigma(x_1, \vec{k}_\perp)|^2. \quad (17)$$

Actually, the root mean square transverse momentum (r.m.s.), $\langle k_\perp^2 \rangle^{1/2}$, should be larger than 300 MeV⁹. Lepage et al. [8], examining the process $\pi^0 \rightarrow \gamma\gamma$, derived yet another constraint on the pion wave function:

$$1 = \frac{f_\pi^2}{12} \int dx_1 \phi(x_1) \Sigma(x_1, \vec{k}_\perp = 0). \quad (18)$$

For the $k_\perp$ -dependence of the wave function we assume a simple Gaussian

$$\Sigma(x_1, \vec{k}_\perp) = 16\pi^2 \beta^2 g(x_1) \exp(-g(x_1) \beta^2 k_\perp^2), \quad (19)$$

$g(x_1)$ being either 1 or $1/x_1 x_2$. In the first case the wave function Ψ_0 factorizes in x_1 and $k_\perp$, which manifestly breaks rotational invariance. This theoretical deficiency is unessential to the purpose of this article. The second, non-factorizing case is obtained from a harmonic oscillator rest frame wave function by equating the energy propagators in both the rest frame and the infinite momentum frame [8]. This particular $k_\perp$ -dependence goes along with a factor $\exp[-\beta^2 m_q^2 / x_1 x_2]$ in the DA. Here, m_q is a constituent quark mass for which we choose a value of 330 MeV. The Gaussian (19) is consistent with the required large- $k_\perp$ behaviour of a soft wave function. For the DAs we try two versions, the asymptotic form $\sim x_1 x_2$ and the CZ form $\sim x_1 x_2 (x_1 - x_2)^2$ [11, 12]. Thus all together we utilize four examples of pion wave functions:

$$\phi_a(x) = A_a x_1 x_2 \quad g = 1 \quad (20a)$$

$$\phi_b(x) = A_b x_1 x_2 (x_1 - x_2)^2 \quad g = 1 \quad (20b)$$

$$\phi_c(x) = A_c x_1 x_2 \exp\left[-\frac{\beta_c^2 m_q^2}{x_1 x_2}\right] \quad g = \frac{1}{x_1 x_2} \quad (20c)$$

$$\phi_d(x) = A_d x_1 x_2 (x_1 - x_2)^2 \exp\left[-\frac{\beta_d^2 m_q^2}{x_1 x_2}\right] \quad g = \frac{1}{x_1 x_2} \quad (20d)$$

⁸The use of the soft part Ψ_0 of the wave function in (2) instead of the full one, entails an insignificant $\mathcal{O}(\alpha_f)$ error.

⁹Lepage et al. [8] have proposed to determine the radius, and in turn the r.m.s. transverse momentum, from the derivative $\partial F_\pi / \partial Q^2$ at $Q^2 = 0$ which can be obtained through eq. (1). Results obtained that way are consistent with our ones.

Example b) emphasizes most the end-point regions, example c) least of all. In principle the DAs depend on the renormalization scale. Asymptotically, that is for $Q \rightarrow \infty$, they all evolve into the asymptotic form (20a), which itself has no evolution. We are going to ignore the evolution in the main part of this article since it is a complication of minor importance. We will however comment below on the effects the evolution may cause. Finally, a remark is in order concerning the DAs (20c) and (20d). Such functions have been proposed by Chernyak and Zhitnitsky [11] on the basis of the two lowest moments of the pion DA as derived from QCD sum rules. The CZ ansatz for the DAs is the subject of considerable controversy in which we do not want to enter ¹⁰. We merely consider (20c) and (20d) as examples of DAs strongly concentrated in the end-point regions.

Our wave functions have one free parameter, the oscillator parameter β , which we fix by requiring specific values for the r.m.s. transverse momentum, namely 350 and 250 MeV. In table 1 we compile the properties of our wave functions, namely the constants A , the oscillator parameters β , the probabilities of the $q\bar{q}$ Fock state $P_{q\bar{q}}$ and the value of the right-hand side of eq. (18), the $\pi \rightarrow \gamma\gamma$ constraint. The value of 350 MeV for the r.m.s. transverse momentum turns out to be a reasonable value on all accounts. The corresponding radius is a little smaller than the measured charge radius of the pion and the constraint (18) is well satisfied. The probabilities of the $q\bar{q}$ Fock state also have plausible values. On the other hand, the value of 250 MeV for the r.m.s. transverse momentum is unrealistic. The radius is too large and (18) is strongly violated. We, however, keep this value for comparison.

Numerical results: The Fourier transform of the $k_\perp$ -dependent part of the wave function reads

$$\hat{\Sigma}(x_1, \vec{b}) = 4\pi \exp\left(-\frac{b^2}{4g(x_1)\beta^2}\right). \quad (21)$$

Li and Sterman [5] assume that the dominant b -dependence of the integrand in eq. (9) arises from the Sudakov factor and that the Gaussian in $\hat{\Sigma}$ can consequently be replaced by 1. In order to examine that assumption, we compare in Fig. 1 the Gaussian with the Sudakov factor [see (13)]. We only display the case $g = 1$ since the other case, $g = 1/x_1x_2$, is very similar. Obviously the intrinsic $k_\perp$ -dependence of the wave function cannot be ignored. For momentum transfers of the order of a few GeV the wave function damps the integrand in (9) more than the Sudakov factor. Only at very large values of Q does the Sudakov factor take over. What is more important at a given value of Q , the Sudakov factor or the wave function depends on the value chosen for Λ_{QCD} and $\langle k_\perp^2 \rangle^{1/2}$. For the favoured values, 200 MeV for Λ_{QCD} and 350 MeV for the r.m.s. transverse momentum, the wave function plays the major rôle in the large b behaviour of the integrand in (9) for Q less than 20 GeV, whereas the Sudakov factor begins to dominate beyond that value, which is well above the experimentally accessible momentum transfer region.

Numerical evaluations of the pion form factor through (9), using the wave functions given in (20), confirm the observations made in Fig. 1. For the two extreme DAs, the one that is most concentrated in the end-point regions, (20b), and the one that is least, (20c), we display the results in Fig. 2. The results for the other two examples, (20a) and (20d), lie in between the two extreme cases. As compared with the hard-scattering result, eq. (6), the inclusion of the Sudakov factor leads to some suppressions as already noticed in [5]. The intrinsic $k_\perp$ -

¹⁰For instance, Braun and Filianov [13], also employing QCD sum rule techniques, found a value of 1.2 ± 0.3 for the pion DA at $x = 1/2$, in clear contradiction with the DAs of the CZ type. It has been pointed out by Radyushkin [3] that the finite size of the vacuum fluctuations had not been considered in [11]. Taking these finite-size effects into account, one finds values for the moments of the pion DA that are much closer to those obtained from the asymptotic DA than the values quoted in [11].

dependence of the wave functions provides additional suppression which amounts to about 50%, in the case (20b), at $Q = 2$ GeV and is still substantial at 7 GeV. As expected the total suppression obtained from both the Sudakov factor and the intrinsic $k_\perp$ -dependence is stronger for the wave function (20b) than for (20c). One may regard the Sudakov factor as a part of the wave function, in fact as its large- b behaviour. In this case one would not consider the product of the two functions, $\exp[-S]$ and $\hat{\Sigma}$. Rather, to an admittedly crude approximation, one would take either the Gaussian in $\hat{\Sigma}$ or the Sudakov factor, whichever is the smaller at a given value of b . Fortunately this recipe yields only minor differences with the first case.

The strong suppression due to the intrinsic $k_\perp$ -dependence of the wave function reveals that the approximation made in the derivation of the hard-scattering formula (6), namely the neglect of the $k_\perp$ -dependence in the hard-scattering amplitude (4), is invalid at small momentum transfer. The reason for this can easily be understood. Using, for instance, a wave function of the type (15,19) with $g = 1$ and, for the sake of the argument, replacing α_s in (4) by an average value, one can convert (3) into

$$Q^2 F_\pi(Q^2) = \frac{2\pi}{3} f_\pi^2 \bar{\alpha}_s C_F \int_0^u dz e^z E_1(z) \int_z^u \frac{dy}{y} \phi(y/u) \phi(z/y), \quad (22)$$

where $u = Q^2 \beta^2 / 2$ and E_1 is the exponential integral. The dominant contribution comes from the upper limit of the z integration. Replacing E_1 by the leading term of its asymptotic expansion

$$z e^z E_1(z) \sim 1 - \frac{1}{z} + \frac{2}{z^2} \mp \dots, \quad (23)$$

one obtains the hard-scattering contribution (6). But the higher-order terms provide substantial corrections to it. The wave function (20a), for instance, leads to

$$Q^2 F_\pi(Q^2) = Q^2 F_\pi^{HSP}(Q^2) \left[1 - 48 \frac{\langle k_\perp^2 \rangle}{Q^2} + \dots \right]. \quad (24)$$

In Fig. 2 we have also shown the available data for the pion form factor at large momentum transfer [9]. This is not meant as a serious comparison between data and theory since, as we mentioned above, the data may suffer from large systematic errors [10]. Nevertheless, it seems to us that the perturbative contribution (9), including the Sudakov corrections and the intrinsic $k_\perp$ -dependence of the wave function, is too small with respect to the data, perhaps by a factor of 2 or so. One may suspect the QCD evolution of the DAs, ignored by us up to now, to be responsible for that discrepancy. Let us estimate whether this explanation holds or not. The evolution equation for the pion DA has been derived in [1]. For our example (20b) it reads

$$\phi_b(x_1, w) = \phi_a(x_1) \left[1 + \frac{2}{3} C_2^{3/2}(x_1 - x_2) \left[\frac{\alpha_s(w^2)}{\alpha_s(\mu_0^2)} \right]^{50/81} \right], \quad (25)$$

where the $C_n^{3/2}(\zeta)$ are the Gegenbauer polynomials, which are the eigensolutions of the evolution equation. According to [6] the evolution parameter w is taken to be $1/b$. Inserting (25) into eq. (9) and using $\mu_0 = 0.5$ GeV [11], we re-evaluate the pion form factor and find that the evolution reduces even further the predictions for F_π (by about 30% for Q between 3 and 4 GeV). Similar results are obtained for the other examples of wave functions, (20c) and (20d). This is in agreement with the findings reported in [12].

Thus we are tempted to suppose that higher-twist contributions, besides the $k_\perp$ -dependence, are quite strong in the few GeV region. Sources of such higher twists are contributions from higher Fock states, $L \neq 0$ components in the valence Fock state wave function or genuine soft

contributions.

An example of genuine soft contributions is provided by the starting point of our considerations, namely the Drell-Yan formula (1). Up to now we have only considered those contributions from (1) which arise from the perturbative tail of the wave functions. The size of the ignored soft contributions until now can easily be estimated for our wave functions (20). One finds

$$F_{\pi}^{soft} = \frac{\pi^2}{3} f_{\pi}^2 \beta^2 \int dx_1 g(x_1) \phi^2(x_1) \exp\left(-g(x_1) \beta^2 x_2^2 Q^2/2\right). \quad (26)$$

The integral is dominated by the region near $x_1 = 1$. Other x_1 regions are strongly damped by the Gaussian. Hence F_{π} sensitively reacts to the behaviour of the DA for $x_1 \rightarrow 1$. Numerical evaluations of (26) for our wave functions (20) show that the examples (20a) and (20c) provide soft contributions of the right magnitude to fill in the gap between the perturbative contribution (9) and the experimental data ¹¹ ($Q^2 F_{\pi}^{soft} \simeq 0.25 \text{ GeV}^2$ at $Q = 2 \text{ GeV}$). As required by the consistency of the entire picture, F_{π}^{soft} decreases faster with increasing Q than the perturbative contribution. The exponential $\exp(-\beta^2 m_q^2/x_1 x_2)$ in the example (20c) turns out to be quite important since it is effective in the end-point regions. It reduces the size of $Q^2 F_{\pi}^{soft}$ substantially and alters its asymptotic behaviour from a $1/Q^2$ behaviour to an exponentially damped decrease; $Q^2 F_{\pi}^{soft}$ becomes equal to the perturbative contribution at $Q \simeq 5 \text{ GeV}$. The soft contributions are also subject to Sudakov corrections. An estimate of these corrections on the basis of eq. (13) reveals, however, that they only amount to a few per cent for the examples (20a) and (20c). The wave functions (20b) and (20d) provide very large soft contributions because of their strong concentration in the end-point regions. Therefore, these examples appear to be unrealistic [see also the discussion after eq. (20)]. Soft contributions of the type (26) have extensively been discussed in [2]. Our results for F_{π}^{soft} are similar in trend, but smaller in size than those presented in [2]. There are three reasons for the differences in the size of the soft contributions: Isgur et al. [2] assume that $P_{q\bar{q}} = 1$, they neglect the exponential $\exp(-\beta^2 m_q^2/x_1 x_2)$ and utilize DAs that are proportional to $\sqrt{x_1 x_2}$ instead of $\sim x_1 x_2$. Strong soft contributions to the pion form factor have also been obtained with QCD sum rules [3]. Finally we mention $L \neq 0$ components of the wave function as another source of higher-twist contributions. For instance, the valence Fock state component of the pion may be expressed by

$$\int \frac{dx_1 d^2 k_{\perp}}{16\pi^3} \frac{1}{\sqrt{2}} (\not{p} + m) \left[\Psi^0(x_1, \vec{k}_{\perp}) + \not{k}_{\perp} \Psi^1(x_1, \vec{k}_{\perp}) \right] \gamma_5, \quad (27)$$

where p denotes the pion's momentum and m its mass. Ralston and Pire [15] consider an even more general ansatz for the valence Fock state wave function of the pion. Thus one may have additional $k_{\perp}$ -dependent contributions to the pion form factor, which are not necessarily small. Perhaps one obtains a better description of the pion form factor with such contributions although at the expense of the introduction of a new, a priori unknown, phenomenological function. Contributions from higher Fock states suffer from the same disadvantage; also in this case new phenomenological functions have to be introduced. Such extensions of the hard-scattering model are certainly not very attractive but are, perhaps, necessary.

Summary: On the basis of our numerical studies we conclude that the intrinsic $k_{\perp}$ -dependence of the wave function has to be taken into account for a reliable quantitative estimate of the

¹¹The Drell-Yan formula (1) has been obtained by a transverse boost to the infinite-momentum frame. Since we know the hadronic wave function only approximately, it may happen that other boost directions provide different results for the pion form factor. For a discussion of such eventual violations of rotational invariance, we refer to [2, 14].

perturbative QCD contribution to the pion form factor. We are aware that this introduces a new phenomenological element into the calculation. The disadvantage is, at least partially, compensated by the fact that the inclusion of the intrinsic $k_{\perp}$ -dependence renders the perturbative contribution even more self-consistent than the Sudakov suppression already does. Applying the criterion of self-consistency as suggested by Li and Sterman [5] (see above), we can conclude that perturbative QCD begins to be self-consistent for Q between 1 and 2 GeV (for $\langle k_{\perp}^2 \rangle^{1/2} = 350$ MeV). The value for Q at which self-consistency sets in depends on the wave function. It is larger for the end-point concentrated wave functions (20b,d) than for the other examples (20a,c). However, the perturbative contribution, although self-consistent, is presumably too small with respect to the data. It thus seems that other contributions (higher twists) also play an important rôle in the few GeV region. We see no argument why our observations should not apply as well to other exclusive observables, such as the nucleon form factor.

References

- [1] G. P. Lepage and S. J. Brodsky, Phys. Rev. D22 (1980) 2157
- [2] N. Isgur and C. H. Llewellyn Smith, Nucl. Phys. B317 (1989) 526
- [3] A. V. Radyushkin, Nucl. Phys. B325 (1991) 141c
- [4] J. Botts and G. Sterman, Nucl. Phys. B325 (1989) 62
- [5] H. Li and G. Sterman, Nucl. Phys. B381 (1992) 129
- [6] H. Li, preprint ITP-SB-92-25, Stony Brook (1992)
- [7] S. D. Drell and T. M. Yan, Phys. Rev. Lett. 24 (1970) 181
- [8] G. P. Lepage, S. J. Brodsky, T. Huang and P. B. Mackenzie, Banff Summer Institute, Particles and Fields 2, p. 83, A. Z. Capri and A. N. Kamal (eds.), 1983
- [9] C. J. Bebek et al., Phys. Rev. D13 (1976) 25 and D17 (1978) 1693
- [10] C. E. Carlson and J. Milana, Phys. Rev. Lett. 65 (1990) 1717
- [11] V. L. Chernyak and A. R. Zhitnitsky, Nucl. Phys. B201 (1982) 492
- [12] T. Huang and Q.-X. Sheng, Z. Phys. C50 (1991) 139
- [13] V. M. Braun and I. E. Filianov, Z. Phys. C44 (1989) 157
- [14] M. Sawicki, Phys. Rev. D46 (1992) 474
- [15] J. P. Ralston and B. Pire, preprint Kansas 5-15-92 (1992)

Table 1: Properties of the pion wave functions (20).

DA	A	$\beta^2[\text{GeV}^{-2}]$	$P_{\bar{q}q}$	R.h.s. of (18)	A	$\beta^2[\text{GeV}^{-2}]$	$P_{\bar{q}q}$	R.h.s. of (18)
a)	6	4.082	0.284	0.947	6	8.0	0.557	1.856
b)	30	4.082	0.338	0.947	30	8.0	0.663	1.856
c)	10.07	0.883	0.320	1.044	16.41	1.787	0.681	2.006
d)	50.05	0.571	0.357	0.983	83.59	1.247	0.721	1.915
$\sqrt{\langle k_T^2 \rangle} = 350 \text{ MeV}$					$\sqrt{\langle k_T^2 \rangle} = 250 \text{ MeV}$			

Figure captions

Fig. 1: The Sudakov factor, evaluated at $x_1 = y_1 = 1/2$, and the Gaussian $\exp(-b^2/2\beta^2)$ [see eq. (21)] as functions of the transverse separation b . The Gaussian is shown for a r.m.s. transverse momentum of 350 MeV (short-dashed line) and 250 MeV (long-dashed line). The Sudakov factor is evaluated for $\Lambda_{QCD} = 200 \text{ MeV}$ (solid lines) and 100 MeV (dotted lines) for $Q_1 = 2 \text{ GeV}$, $Q_2 = 5 \text{ GeV}$ and $Q_3 = 20 \text{ GeV}$ ($x_1 = x_2 = 1/2$).

Fig. 2a: The pion form factor as a function of Q^2 evaluated with the wave function (20b) and $\Lambda_{QCD} = 200 \text{ MeV}$. The dash-dotted line is obtained from the hard-scattering formula with an α_s cut-off at 0.5 and the dashed line from (9) ignoring the intrinsic $k_\perp$ -dependence. The solid and the dotted lines represent the complete result obtained from (9) taking into account both the Sudakov factor and the intrinsic $k_\perp$ -dependence for $\langle k_\perp^2 \rangle = 350 \text{ MeV}$ and 250 MeV, respectively. Data are taken from [9] ($\circ$ 1976, $\bullet$ 1978).

Fig. 2b: As Fig. 2a but using the wave function (20c). Note the modified scale of the abscissa.




Fig. 1




Fig. 2a




Fig. 2b